



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{2r^3}{a\beta\gamma} + \left(\frac{1}{a^2} + \frac{1}{\beta^2} + \frac{1}{\gamma^2}\right)r^2 = 1, \text{ or } r^3 + \frac{1}{2}\left(\frac{a\beta}{\delta} + \frac{\beta\gamma}{a} + \frac{\gamma a}{\beta}\right)r^2 = \frac{1}{2}a\beta\gamma.$$

Example. Let $a=275$, $\beta=325$, $\gamma=429$; then $r^3 + 539\frac{1}{6}r^2 - 19170937\frac{1}{2} = 0$; whence $(r-165)(r^2 + 704\frac{1}{6}r + 116187\frac{1}{2}) = 0$.

$\therefore r=165$. Also, $r=-137.97+$ and $r=842.13+$.

CALCULUS.

140. Proposed by C. C. BEBOUT, Professor of Mathematics, Elgin High School, Elgin, Ill.

A pole two inches in diameter is set vertically in a level plat of ground. At a point ten feet from the ground a string is attached. A man holds the other end of the string and walks about the pole keeping the string stretched taut, and his hand at a constant distance of four feet from the ground, till the string is all wound upon the pole. If string is ten feet long, how far has his hand moved in the operation?

Solution by G. B. M. ZERR, A.M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let n be the number of times the string 10 feet in length will encircle the cylinder in a height of 6 feet. Circumference of cylinder $= \frac{1}{6}\pi$ feet, x = distance between successive portions of the string on the same generating line.

Then $\sqrt{[(\frac{1}{6}\pi)^2 + x^2]} = 10/n$ also $nx=6$.

$$\therefore n \sqrt{\left(\frac{\pi}{6}\right)^2 + \left(\frac{6}{n}\right)^2} = 10 \text{ or } \frac{\pi^2 n^2}{36} = 64. \therefore n = \frac{48}{\pi}, x = \frac{1}{8}\pi.$$

The distance moved by the hand is given on page 319, No. 9, Vol. I, of the MONTHLY, and is there worked out in full. It is $s=2\pi^2 r^2 m^2$ where r is the radius of the cylinder and m is the number of times the string is wound around the cylinder or the same as the value of n above. $r=\frac{1}{12}$ feet, $m=n=48/\pi$.

$\therefore s=32$ feet.

Also solved by T. T. DAVIS.

141. Proposed by B. F. FINKEL, A.M., M.Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

The curve $r^n = a^n \sin n\theta$ rolls along a straight line. Show that the intrinsic equation to the evolute of the locus of the pole is $s^n = a^m [1 + 1/n]^n \sin \psi$. [Edward's *Differential Calculus*, page 502.

Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

The distance from the point of tangency to the pole is normal to the path of the pole. Let this distance, which is a radius vector of the rolling curve, be r , and the perpendicular from the pole upon the fixed line be p .

Taking the fixed line as the axis of x , p is an ordinate of the pole, and r is the length of normal and is expressed by